

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

REPLACEMENT DRAWINGS

1/17



Mechanical Shearing of CEPH 1347-2 gDNA using the TECAN




FIG. 1A

Boiling or DNase I-digestion of CEPH 1347-2 gDNA




FIG. 1B

REPLACEMENT DRAWINGS

2/17

Effect of gDNA Concentration and Boiling Duration on Range of Fragment Size Generated




FIG. 2

REPLACEMENT DRAWINGS

3/17

Fragment Sizes Generated by Boiling of
Different Sources of gDNA




FIG. 3

REPLACEMENT DRAWINGS

4/17





FIG. 4

Boiled gDNA can Participate in OLA-PCR Reactions



Lane 2: 25 bp marker
 Lane 3: DNase I-digested CEPH 1347-2 gDNA
 Lane 4: CEPH 1347-2 gDNA, boiled 15 min.
 Lane 5: CEPH 1347-2 gDNA, boiled 30 min.
 Lane 6: Ligation control (no gDNA)
 Lane 7: DNase I-digested NA-12565 gDNA
 Lane 8: NA-12565 gDNA, boiled 15 min.
 Lane 9: NA-12565 gDNA, boiled 30 min.
 Lane 10: Ligation control (no gDNA)

FIG. 5A

REPLACEMENT DRAWINGS

5/17




FIG. 5B

6/17

OLA-PCR Products Formed using Boiled gDNA Binds to PE-27 Arrays




FIG. 6A

REPLACEMENT DRAWINGS

7/17




FIG. 6B

REPLACEMENT DRAWINGS

8/17

OLA-PCR Products Formed using Boiled gDNA Binds to PE-27 Arrays




FIG. 6C

REPLACEMENT DRAWINGS

9/17

OLA-PCR Products Formed using Boiled gDNA Binds to PE-27 Arrays




FIG. 6D

REPLACEMENT DRAWINGS

10/17

OLA-PCR Products Formed using Boiled gDNA Binds to PE-27 Arrays




FIG. 6E

REPLACEMENT DRAWINGS

11/17

OLA-PCR Products Formed using Boiled gDNA Binds to PE-27 Arrays




FIG. 6F

REPLACEMENT DRAWINGS

12/17




FIG. 7A

13/17

OLA-PCR Products Generated using Intact or Boiled gDNA Binds to PE-27 Arrays




FIG. 7B

REPLACEMENT DRAWINGS

14/17

OLA-PCR Products Generated using Intact or Boiled gDNA Binds to PE-27 Arrays




FIG. 7C

REPLACEMENT DRAWINGS

15/17




FIG. 7D

REPLACEMENT DRAWINGS

16/17

OLA-PCR Products Generated using Intact or Boiled gDNA Binds to PE-27 Arrays




FIG. 7E

REPLACEMENT DRAWINGS

17/17




FIG. 7F